Invisible Forces behind G-Protein Coupled Receptors



February 25, 2026

Abstract:

G-protein coupled receptors (GPCRs) are membrane proteins essential for cellular communication and are targeted by over one-third of modern pharmaceuticals. Rhodopsin, a light-sensitive GPCR in the retina, serves as a model for studying GPCR activation because it transitions between distinct functional states, including the inactive Meta I and active Meta II intermediates. Although GPCRs are widely studied, many experiments are conducted under artificial conditions that do not replicate complex environments present in biological systems. While previous studies have demonstrated that lipid composition and surrounding water molecules influence rhodopsin activation, less attention has been given to how ionic strength regulates the Meta I/Meta II equilibrium under physiologically relevant conditions.

In this study, rhodopsin samples were analyzed under identical experimental conditions while ionic strength was varied across 0.5×, 1.0× (control), and 2.0× salt concentrations. UV-visible spectroscopy monitored conformational changes, with absorbance features near 500 nm, 487 nm, and 380 nm corresponding to the dark state, Meta I, and Meta II, respectively. Difference spectra and replicate measurements were used to compare intermediate states and ensure reliability.

Results showed consistent characteristic spectral features across conditions, while Meta II-associated absorbance decreased under both reduced and elevated ionic strength relative to control. These findings suggest ionic strength modulates rhodopsin activation through electrostatic screening of charged residues, altering conformational equilibrium without changing structural identity. Overall, this work highlights the importance of ionic environments in regulating GPCR function and supports development of physiologically relevant models for studying membrane protein activation and GPCR-targeted drug design.

Acknowledgements:

I would like to sincerely thank Dr. Michael F. Brown, my Principal Investigator, and Zachary T. Bachler and Evelyn W. Cheng, my lab mentors, for their guidance, support, and expertise throughout this project. Their insights and encouragement were invaluable in shaping the direction of my research and in helping me navigate experimental challenges. I also thank STAR Lab for providing the opportunity to conduct this study, as well as for the resources and mentorship that made this work possible.

Table of Contents

Introduction ……..……..…….. 1

Research Question ……..…….. 2

Hypothesis ……..……..…….. 2

Methods ……..……..…….. 3

Results ……..……..……..…….. 4

Discussion ……..……..…….. 8

Conclusion ……..……..……..……..…… 9

References ……..……..……..…… 11

Introduction:

Membrane proteins are critical for nearly every aspect of cellular communication and signaling. Among them, G-protein coupled receptors (GPCRs) form one of the largest and most physiologically significant protein families, responsible for detecting external signals such as light, odorants, and hormones, and transmitting them into cellular responses [Rasmussen et al., 2019]. Because GPCRs regulate many essential biological processes, they serve as targets for more than one-third of all modern pharmaceuticals [Hauser et al., 2017]. However, despite their importance, understanding the mechanisms that govern GPCR activation and function remains challenging, largely due to the difficulty of studying these proteins within their complex and dynamic membrane environments. Traditional studies often isolate membrane proteins in artificial detergent micelles, which fail to accurately represent the natural conditions found in biological membranes. As a result, important aspects of lipid-protein interactions, stability, and signaling dynamics may be overlooked.

A well-established model for understanding GPCR structure and activation is the membrane protein rhodopsin, a light-sensitive GPCR located in the photoreceptor cells of the retina. In dim light conditions, rhodopsin enables vision by initiating a signaling cascade that converts photons into neural signals. Rhodopsin transitions between multiple conformational states during activation, with the Meta I state representing an intermediate inactive form and the Meta II state representing the active conformation capable of triggering downstream signaling. The equilibrium between Meta I and Meta II is sensitive to environmental conditions, and shifts in this balance directly affect rhodopsin function. While decades of biochemical and structural research have provided a foundational understanding of the activation of rhodopsin, recent studies emphasize that its surrounding environment — particularly the water molecules and lipid membrane composition — plays a critical role in regulating its behavior [Brown et al., 2009; Bachler & Brown, 2024].

Earlier investigations have demonstrated that the lipid composition surrounding rhodopsin can influence both its structural conformation and activation efficiency. Using solid-state nuclear magnetic resonance (NMR), it was found that the lipid composition and membrane packing density directly modulate rhodopsin's conformational changes, suggesting that membrane properties can regulate receptor function [Brown et al., 2009]. More recent work by Bachler and Brown [2024] revealed that distinct populations of water molecules surrounding rhodopsin exhibit distinct effects on protein activation — some remaining tightly bound to the

protein surface, while others facilitate larger conformational shifts. Together, these findings indicate that the surrounding water and lipid environments jointly orchestrate how rhodopsin — and potentially other GPCRs — transitions between active and inactive states.

In addition to lipids and water, the chemical environment surrounding rhodopsin — particularly the composition of the buffer in which it is studied — plays a major role in shaping its structural dynamics. Biological buffers are designed to maintain stable pH, control ionic strength, and preserve protein stability during experiments. These factors are essential for membrane proteins like rhodopsin, whose activation involves subtle shifts in protonation states and electrostatic interactions. Because many studies rely on simplified or non-physiological buffer conditions, key environmental influences such as ionic strength may be unintentionally altered or overlooked. As a result, understanding how buffer composition modulates rhodopsin's conformational equilibrium is crucial for accurately interpreting its activation behavior.

Despite these advances, less attention has been given to the influence of ionic strength on rhodopsin activation. Because salt concentration directly affects electrostatic interactions and protein stability, variations in ionic strength may serve as a critical but underexplored regulator of GPCR function. This study addresses this gap by investigating how ionic strength influences rhodopsin's structure, activation, and function in membrane environments that more closely mimic native biological systems. By integrating realistic membrane conditions, this research aligns with the Brown Lab's broader mission to uncover how a protein's physical surroundings affect its behavior and signaling capabilities. The findings may provide generalizable insights into the environmental factors that modulate GPCR function, ultimately aiding in the development of more effective and physiologically informed drug designs.

Research Question:

How does varying salt concentrations in natural biological membranes regulate the structure, activation, and function of rhodopsin?

Hypothesis:

Increasing the salt concentration will shift the Meta I and Meta II equilibrium to favor the Meta II (active) state. This is because higher ionic strength alters membrane surface charge and screens electrostatic interactions, making it easier for rhodopsin to undergo the structural change required for activation.

Methods:

UV-visible (UV-Vis) spectroscopy was used to monitor conformational changes in rhodopsin under varying experimental conditions. UV-Vis spectroscopy measures light absorbance across a range of wavelengths, providing a sensitive, non-invasive method for detecting structural changes in light-activated proteins. Rhodopsin exhibits distinct absorption spectra in its major functional states — the dark state absorbing at 500 nm, Meta I at 487 nm, and Meta II at 380 nm — allowing transitions between these states to be tracked through changes in absorbance peaks. Difference spectra were generated by subtracting the dark-state spectrum from the light-activated spectrum, highlighting wavelengths where changes occur and making it easier to detect subtle shifts in absorbance.

Rhodopsin samples were prepared under controlled conditions in bis-tris propane (BTP) buffer, a conjugate acid-base system used to maintain stable pH. Ionic conditions were systematically varied by adjusting the concentration of salts (NaCl and $MgCl_2$) added to the buffer. All other experimental parameters, including pH, temperature, and overall chemical environment, were held constant. UV-Vis absorption spectra were recorded for each condition before and after light activation to allow direct comparison across salt concentrations. Dark spectra were collected first to establish a baseline corresponding to rhodopsin's inactive state. These baseline measurements were compared with light-activated spectra to determine absorbance shifts and quantify transitions between Meta I and Meta II.

Spectral data were analyzed by examining absorbance changes at key wavelengths associated with rhodopsin's functional states. Difference spectra were generated to quantify shifts indicative of transitions between Meta I and Meta II. Absorbance values were averaged across replicate trials to improve reliability. When multiple conditions were compared, appropriate statistical analyses, such as t-tests or ANOVA, were planned to determine whether observed differences were significant.

Potential risks associated with this project were minimal. Some experimental conditions involved the use of concentrated hydrochloric acid to adjust pH. Hydrochloric acid is corrosive; appropriate safety precautions, including gloves, goggles, and lab coats, were used. All chemical handling followed established laboratory safety protocols.

Results:

UV-visible absorbance spectroscopy was used to compare rhodopsin spectral profiles across three experimental conditions corresponding to 0.5× control concentration (0.5), control concentration (1.0), and 2× control concentration (2.0). Representative samples were selected from each of the experimental conditions.

Distinct absorbance features correspond to established conformational states of rhodopsin. The inactive dark-state form exhibits a characteristic absorbance maximum near 500 nm. Absorbance near approximately 487 nm is associated with the Meta I intermediate, while increased absorbance around 380 nm corresponds to formation of the Meta II state. Difference spectra were generated to quantify the relative shifts in Meta I/Meta II equilibrium under varying salt concentrations.



Fig. 1. UV-visible absorbance spectrum after light activation of rhodopsin under control (1.0 x) ionic strength conditions.

The control condition (1.0x) displays a clear peak around 380 nm, indicating the Meta II intermediate. Absorbance decreases toward longer wavelengths, with a shoulder in the 487 nm region and lower absorbance approaching 500 nm, indicating the loss of the dark state. This spectrum serves as the baseline profile against which the other conditions were evaluated.



Fig. 2. UV-visible absorbance spectrum after light activation of rhodopsin at 0.5× control ionic strength conditions.

In the 0.5× condition (0.5), the overall spectral shape remains similar to the control spectrum, with the 380 nm peak still present. However, absorbance values across the wavelength range are reduced relative to the control. Absorbance near 380 nm is comparatively lower and the feature near 487 nm is less pronounced.



Fig. 3. UV-visible absorbance spectrum after light activation of rhodopsin at 2× control ionic strength conditions.

The 2× condition (2.0) exhibits similar absorbance values across the measured wavelengths when compared with the 0.5× conditions. The overall spectral shape remains constant. However, absorbance values across the wavelength range are reduced relative to the control. Absorbance near 380 nm is comparatively lower and the feature near 487 nm is less pronounced.

Across all three conditions, the locations of major absorbance features remain consistent, with variation observed in absorbance intensity rather than wavelength shifts. The spectra therefore differ mainly in relative magnitude while preserving the characteristic rhodopsin absorbance profile.

Discussion:

Rhodopsin activation is a fundamental process in visual phototransduction, in which structural transitions between intermediates regulate downstream signaling. Since rhodopsin serves as a model GPCR, its activation process can also be applied to other GPCRs. This study investigated how changes in salt concentration influence the equilibrium between rhodopsin intermediates, particularly the Meta I and Meta II states. Because electrostatic interactions play a critical role in stabilizing rhodopsin structure, altering ionic strength provides a way to test how environmental conditions affect these conformational transitions.

Across experimental conditions, the absorbance spectra retained characteristic rhodopsin features, indicating that overall protein structure remained intact. However, relative absorbance intensities varied with salt concentration, suggesting shifts in the equilibrium distribution of conformational states rather than changes in spectral identity. Compared with the control condition, both decreased (0.5×) and increased (2×) salt concentrations showed reduced absorbance associated with Meta II formation, indicating that ionic strength influences the efficiency or stability of rhodopsin activation.

These observations can be interpreted through known molecular interactions within rhodopsin. A key structural feature involves an electrostatic interaction between the amino acids glutamate at position 134 (Glu134), which normally carries a negative charge, and arginine at position 135 (Arg135), which is positively charged. In the Meta I state, these residues form a stabilizing interaction. Transition to the Meta II state occurs when Glu134 becomes protonated, neutralizing its charge and weakening this interaction, allowing the protein to undergo conformational rearrangement associated with activation. Increasing salt concentration introduces additional ions into the solution that can interact with charged residues. These ions effectively weaken electrostatic attractions by forming alternative interactions with Glu134 and Arg135, reducing the strength of their mutual attraction. As a result, the balance of forces governing conformational stability is altered.

The reduced Meta II–associated absorbance observed under altered salt conditions suggests that ionic screening may interfere with the precise electrostatic changes required for efficient Meta II stabilization. Rather than promoting activation, excess ions may compete with intramolecular interactions, preventing the structural rearrangements from fully stabilizing the active conformation. Similarly, lower ionic strength may reduce overall electrostatic stabilization within the protein environment, also limiting efficient formation of Meta II. Together, these findings suggest that rhodopsin activation depends on an optimal ionic environment that preserves critical internal charge interactions while still allowing protonation-driven conformational change.

These results are consistent with broader studies of G-protein coupled receptors (GPCRs), in which activation often depends on finely balanced electrostatic networks and protonation events. Previous work has shown that ionic strength can modulate protein conformational equilibria by altering long-range charge interactions, supporting the interpretation that salt-dependent screening influences rhodopsin intermediate stability.

Several limitations should be considered when interpreting these findings. First, UV–visible spectroscopy measures ensemble absorbance rather than direct structural changes, meaning that conformational assignments rely on established spectral correlations rather than direct observation. Second, variations in absorbance magnitude could partially reflect experimental factors such as pipetting variability, protein concentration differences, or instrumental noise. Additionally, only three salt conditions were tested, limiting the ability to determine whether the relationship between ionic strength and Meta II formation is linear or follows a more complex trend.

Future research could expand this work by testing a broader range of ionic strengths or comparing different ion species to evaluate specific ion effects.

Conclusion:

This study examined how salt concentration influences rhodopsin conformational equilibria using UV-visible absorbance spectroscopy. The results indicate that while characteristic spectral features remain unchanged, variations in ionic strength alter absorbance intensity associated with the Meta II intermediate, suggesting modulation of rhodopsin activation efficiency rather than structural identity. These findings support a model in which electrostatic

interactions — particularly between Glu134 and Arg135 — are sensitive to ionic screening effects in solution.

Overall, the results suggest that rhodopsin activation requires a balanced electrostatic environment, highlighting the importance of intramolecular charge interactions in GPCR function. By demonstrating how external ionic conditions influence conformational equilibrium, this work contributes to a broader understanding of protein activation mechanisms and raises new questions about how cellular environments regulate photoreceptor signaling dynamics.

References

Bachler, Z. T., & Brown, M. F. (2024). Hidden water's influence on rhodopsin activation. *Biophys. J.*, 123(24), 4167–4179.

Brown, M. F. (2012). Curvature forces in membrane lipid–protein interactions. *Biochem.*, 51(49), 9782–9795.

Brown, M. F., Martínez-Mayorga, K., Struts, A. V., & Nakanishi, K. (2009). Retinal conformation and dynamics in activation of rhodopsin illuminated by solid-state ²H NMR spectroscopy. *Photochem. Photobiol.*, 85(2), 442–453.

Fried, S. D. E., Lewis, J. W., Szundi, I., Martínez-Mayorga, K., Mahalingam, M., Vogel, R., Kliger, D. S., & Brown, M. F. (2021). Membrane curvature revisited—the archetype of rhodopsin studied by time-resolved electronic spectroscopy. *Biophys. J.*, 120(3), 440–452.

Hauser AS, Attwood MM, Rask-Andersen M, Schiöth HB, Gloriam DE. Trends in GPCR drug discovery: new agents, targets and indications. *Nat Rev Drug Discov.* 2017;16(12):829-842.

Rasmussen SGF, DeVree BT, Zou Y, et al. Crystal structure of the β2 adrenergic receptor–Gs protein complex. *Nature*. 2011;477(7365):549-555.

Struts, A. V., Barmasov, A. V., Fried, S. D. E., Hewage, K. S. K., Perera, S. M. D. C., & Brown, M. F. (2024). Osmotic stress studies of G-protein-coupled receptor rhodopsin activation. *Biophys. J., Chem.*, 304, 107112.